

52
P93c
no. 8

Nat. Hist.

NOV 16 1926

The Cambrian-Ordovician Stratigraphic Column in Southeastern Newfoundland

(PRINCETON UNIVERSITY CONTRIBUTION TO THE GEOLOGY OF
NEWFOUNDLAND, No. 8)

BY

B. F. HOWELL

Reprinted from *The Canadian Field-Naturalist*, Vol. XL, No. 3,
pp. 52-57, March, 1926

GEOLOGY

557
P93c
no. 8

Nat. Hist.

THE LIBRARY OF THE
NOV 16 1926
UNIVERSITY OF ILLINOIS



(Reprinted from *The Canadian Field-Naturalist*, Vol. XL, No. 3,
pp. 52-57, March, 1926)




OTTAWA
THE GRAPHIC PUBLISHERS

16 Nov 26 dir. g. v. 8 1926 cont. E.H.

THE CAMBRIAN-ORDOVICIAN STRATIGRAPHIC COLUMN OF SOUTHEASTERN NEWFOUNDLAND

By BEN. F. HOWELL

INTRODUCTION



THE sedimentary rocks, ranging in age from Lower Cambrian to late Lower or early Middle Ordovician, which have been found in southeastern Newfoundland, are of more than ordinary interest to stratigraphers and palaeontologists because of the abundance, variety, and excellent preservation of their fossils, and of the remarkable way in which these fossils resemble those found in beds of corresponding age in northwestern Europe. Much work will have to be done on many of the Newfoundland beds and fossils before we shall be in a position to give a detailed account of their exact sedimentary and faunal succession, detailed geographic distribution, and local development; but a brief summary of our present knowledge may perhaps be appropriately presented now.

Much of the information on which this paper is based was gathered by members of the three geological expeditions from Princeton University which worked in southeastern Newfoundland in 1912, 1913 and 1914; and the late Professor Gilbert van Ingen, who organized and led these field parties, would have been far more competent than the writer to describe the early Paleozoic rocks of that region. In acceding to the request of the editors to undertake this task, the writer has been actuated by a desire to make available to the scientific world some of the many facts which Professor van Ingen had gathered together, but which he was unable to publish because of ill health and the pressure of other duties. The writer would also express his indebtedness to the late Messrs. Alexander Murray and James P. Howley, of the Newfoundland Geological Survey,

who gave us the first detailed account of the sedimentary rocks of southeastern Newfoundland; to the late Dr. G. F. Matthew and to Dr. Charles D. Walcott, who added much to our knowledge of the stratigraphy and palaeontology of the beds; to Dr. Albert O. Hayes who, with Professor van Ingen, first worked out the detailed stratigraphy of the Ordovician section in the Conception Bay district; and to Professor Nelson C. Dale, who described the Cambrian manganese beds about Conception Bay and studied the Cambrian rocks along the shores of Fortune Bay. The Cambrian fossils referred to have been identified by the writer. The identifications of the Ordovician ones are by Professor van Ingen.

THE SECTION

The Lower Paleozoic sediments of southeastern Newfoundland are shales and sandstones, with a few limestones, limy shales and limy nodules in the Lower Cambrian and the overlying Paradoxides beds, and with limy concretions in the Upper Cambrian, and a sandy limestone in the succeeding Tremadocian. Their character, faunas, and order of superposition are indicated in Table 1. Their total thickness has been estimated by Professor van Ingen at about 10,000 feet in the Conception Bay region. As far as we can judge, with our present knowledge, they appear to be naturally separable on lithological and palaeontological characters into five main divisions—the red and green shales and limestones with the *Callavia*, *Protolenus* and *Catadoxides* faunas, at the bottom of the column; the gray and black shales and thin gray limestones holding the *Paradoxides* faunas (the lower beds of this division are some-

times red); the gray and black shales and gray sandstones of the Upper Cambrian, containing the *Olepus* fauna; the gray and brown shales and single sandy limestone of the Tremadocian, with *Bryograptus*, *Shumardia*, *Parabolina*, *Princetonia*, and *Niobe*; and the sandstones, shales, and hematite beds of the Arenigian (and possibly Llandeilan) containing *Sphaerobolus*, *Didymograptus*, *Synomalonotus*, *Schizocrania* and *Westonia*.

These beds occur as widely separated erosion remnants of what was probably, during and following the time of deposition of the later beds at least, a more or less continuous sheet of sediments. Because they have been longest subjected to erosion, the more recent of these beds have been more nearly completely removed, while the most ancient ones have been more extensively preserved.

The Lower Cambrian beds are known to outcrop at Topsail, Manuels, Long Pond Brook, Duff's, and intermediate points on the southeastern side of Conception Bay; at Chapel's Cove, Bacon Cove, Collier's Bay, and Brigus Bay on the western side of Conception Bay; on the western and northern shores of Random Island, and for a considerable distance on the northern shore of Smith Sound, Trinity Bay; and near Little Danzig Cove, which lies southwest of Fortune, on the southern shore of Fortune Bay.

The shales and thin limestones of the Newfoundland series, the Paradoxides Beds, have their best exposure in the gorge of Manuel's Brook, on the southeastern side of Conception Bay, some fifteen miles west of St. John's. This is a locality famous in the annals of North American stratigraphy, and of Cambrian geology in general, as the place where Dr. Walcott, in 1888, found *Paradoxides* beds and "*Olenellus*" beds in the same section—the first time that they had been recognized in America in a single section, and the first time that they had been found anywhere in the world so well developed and richly fossiliferous, and with their stratigraphic positions so clearly shown that their relative ages could be determined beyond question—and was able to demonstrate to everyone's satisfaction that the *Paradoxides* faunas were younger than the *Olenellus* faunas, and not older, as had been commonly believed up to that time. The Newfoundland beds are also well exposed at Topsail, Chamberlin's Brook, Long Pond, and Kelligrew Brook on the southeastern side of Conception Bay; at Chapel's Cove and on the south side of Brigus Head on the west side of Conception Bay; at Chapple Arm and Island Cove, near the head of Trinity Bay; on the west and north shores of Random Island and the north shore of Smith Sound, Trinity Bay (where the lower beds of the series are red shales and limestones); at Branch, St. Mary's Bay; near Fortune on the south side

of Fortune Bay, near Boxey Point on the western shore of the same bay; and on Little Miquelon Island, which is not a part of Newfoundland, but which belongs geologically with that island, as it lies near it geographically.

The Upper Cambrian, "Olenian", Elliott Cove series, forms the part of Random Island which lies southeast of the island's northwestern corner. It is supposed to underlie the mainland on the other side of Smith Sound, to the northeast, though the island along that shore of the Sound is covered with glacial deposits for a considerable distance, and the beds are not known to be well exposed there. The series is also present on the southeastern side of Conception Bay, where it is well exposed in the gorge of Manuels Brook, and on the shore of the bay at Topsail Head. It will probably be found to outcrop also in the valleys of Kelligrew Brook and Seal Cove Brook, southwest of Manuels.

The beds of the Clarendville series, which are of Tremadocian age and older, although they presumably underlie the southeastern part of Conception Bay, are not known to outcrop as a whole above sea-level there, and have been found exposed only about Trinity Bay, where they form the northwestern part of Random Island and a part of the mainland at the western end of Smith Sound, the highest bed of the series, the Riders Brook limestone, occurring only in the latter locality.

The shales and sandstones of the upper division—the Bell Island and Wabana beds, of Arenigian, Lower Ordovician (and perhaps Llandeilan) age—are known only in the Conception Bay basin, and are exposed above sea level only on the three islands, Kelly, Little Bell, and Big Bell Islands, which lie out in that bay, though they must underlie much of the bay, the hematite beds of the Wabana series having been mined for several miles out beneath the sea-bottom northwest of Great Bell Island. The beds which are considered to be possibly of Llandeilan age are exposed on the northwestern side of that island.

In all this great stratigraphic section, from the base of the Cambrian to the uppermost Ordovician beds found, there is no known angular unconformity. Indications that deposition or upward movements of the sea-bottom brought the bottom almost or quite to sea-level have been found at one or more horizons in each of the five stratigraphic divisions described above; but in no case has there been noticed clear evidence of any considerable deformation or subaerial erosion of strata indicating an undoubted stratigraphic break of any importance, though periods of almost complete cessation of deposition seem to have occurred a number of times, and may prove on further

examination to have continued, in some instances, for considerable lengths of time. Concealment of some of the later beds beneath the waters of Conception Bay, between the mainland and Kelly, Little Bell, and Great Bell Islands, and folding of others of the upper beds in Random Island, together with our lack of detailed knowledge of many of the sections on the southern coast-line of southeastern Newfoundland, make it impossible for us to speak more definitely at present about the number or importance of the breaks in this stratigraphic column.

Fossils are common, in some beds at least, in each of the five groups of strata described above. They have not been found in the oldest beds of the Lower Cambrian; they occur in the greatest abundance and variety in the Paradoxidian Newfoundland shales and thin limestones; they are very abundant in some beds of the Upper Cambrian, the Tremadocian, and the Arenigian, but appear to be absent from others. They are sufficiently abundant and well enough distributed in all five divisions to enable us to determine the ages not only of the divisions as a whole, but of most, or all, of the principal parts of each division. They are, too, in many cases so similar to forms occurring in New Brunswick and Cape Breton, on the one side, and Great Britain and Scandinavia, on the other, that we can make close correlations of many of the zones with the corresponding horizons in those regions.

CORRELATION

The correlations of the strata of southeastern Newfoundland with those of the Canadian Maritime Provinces and northwestern Europe are shown in Table 2. An examination of this table will reveal the fact that the zones of the Lower and Middle Cambrian series of southeastern Newfoundland can, in most cases, be accurately identified with zones in southeastern Canada and northwestern Europe, but that the later zones of Newfoundland, while comparable in a general

way with those of New Brunswick, Cape Breton, Great Britain and Scandinavia, cannot be correlated so closely. The main reason for our inability to make closer comparisons between these later beds is undoubtedly our lack of a more detailed knowledge of the Newfoundland faunas—a handicap which will be largely removed when the study of Professor van Ingen's collections, which are at Princeton, has been completed. Further studies of collections already made will probably disclose the age of the shales lying between the *Paradoxides davidis* beds and the *Agnostus pisiformis* beds in Newfoundland, some of which shales may prove to belong in the *Paradoxides forchhammeri* zone. The exact relations of the *Catadoxides* Zone to the *Protolenus* Zone in southeastern Newfoundland is also not well understood. The *Catadoxides* Zone contains an undescribed fauna, which may prove similar to the fauna of the beds lying between the known *Protolenus* beds and the lowest known *Paradoxides* beds of Shropshire, England.

As far as they have yet been studied, the Cambrian-Ordovician faunas of southeastern Newfoundland have proved to be most remarkably similar to the faunas found in strata of the same age in Europe. Further study will undoubtedly add much to our knowledge of the Newfoundland faunas, and will probably enable us not only to recognize a larger number of faunal zones and make further detailed correlations with European and continental North American zones, but will surely disclose the presence of many more species, some of which will resemble closely, or be identical with, European forms, while others will be new or will be forms already known in North America. On the whole, there is good reason to believe that an increased knowledge of the Newfoundland fossils will make this interesting correspondence between the ancient life of the two sides of the Atlantic even more striking that it is now known to be.

BIBLIOGRAPHY OF THE EARLY PALEOZOIC STRATIGRAPHY AND PALEONTOLOGY OF SOUTHEASTERN NEWFOUNDLAND

- Billings, E. 1865. Palaeozoic Fossils, vol. I, Geol. Surv. Canada, 1865.
 Billings, E. 1872. On some new fossils from the Primordial rocks of Newfoundland. The Canadian Naturalist, New Series, vol. 6, 1872, p. 465-479.
 Billings, E. 1874. Palaeozoic Fossils, vol. II, part I, Geol. Surv. Canada, Montreal, 1874.

- Dale, N. C. 1914. Manganese deposits of Conception and Trinity Bays, Newfoundland, Bull. Geol. Soc. Amer., vol. 25, 1914, p. 73, 74
 Dale, N. C. 1915. The Cambrian manganese deposits of Conception and Trinity Bays, Newfoundland. Princeton University Contribution to the Geology of Newfoundland, No. 2. Proc. Amer. Phil. Soc., vol. 54, 1915, p. 317-456.
 Hayes, A. O. 1914. Geology of the Wabana Iron Ore of Newfoundland. Bull. Geol. Soc. America, vol. 25, 1914, p. 74.
 Hayes, A. O. 1915. Wabana Iron Ore of Newfoundland. Geol. Surv. Canada, Memoir 78,

No. 66, Geological Series, 1915.

Howell, B. F. 1920. Correlation of the Middle Cambrian of Newfoundland and Great Britain. Bull. Geol. Soc. Amer., vol. 31, No. 1, 1920, p. 214.

Howell, B. F. 1920. The Middle Cambrian beds at Manuels, Newfoundland, and their relations. Science, New Series, vol. 51, No. 1330, 1920, p. 644.

Howell, B. F. 1925. The faunas of the Cambrian Paradoxides beds at Manuels, Newfound-

land. Princeton University contribution to the Geology of Newfoundland, No. 7. Bulletins of American Paleontology, vol. 11, No. 43; 1925.

Howley, J. P. The Taconic of Eastern Newfoundland. American Geologist, vol. 4, 1889, p. 121-125.

Howley, J. P. 1907. Geological map of Newfoundland, compiled from the most recent and authentic sources, the coast line corrected to date from the admiralty surveys, the interior chiefly

TABLE I.

CAMBRIAN-ORDOVICIAN STRATIGRAPHIC COLUMN OF SOUTHEASTERN NEWFOUNDLAND

SYSTEM	SERIES	ZONE	NATURE OF BEDS	CHARACTERISTIC FOSSILS
	Llandeilan?	<i>Lingula fraseri</i> Zone	Black shale, sometimes phosphatic.	<i>Lingula fraseri</i> <i>Schizocrania striata</i>
	Middle and Upper Arenigian, Wabana Series	Beds with <i>Didymograptus</i> and <i>Synhomalonotus</i>	Black and gray shales, sometimes phosphatic and pyritiferous, gray sandstones, and red oolitic hematite.	<i>Synhomalonotus chambersi</i> <i>Hemigraspis cantleyi</i> <i>Lingula leseuri</i> <i>Schizocrania hayesi</i> <i>Didymograptus nitidus</i>
ORDOVICIAN	Lower Arenigian, Bell Island Series	Beds with <i>Obolus Lingulobolus</i> , and <i>Sphaerobolus</i> .	Brown and gray sandstones and shales, white sandstone, and, at the top, red oolitic hematite.	<i>Obolus burrowsi</i> <i>Sphaerobolus fimbriatus</i> <i>Lingulobolus affinis</i> <i>Lingula hawkei</i> <i>Lingula murrayi</i> <i>Lingula howleyi</i> <i>Lingulella billingsi</i> <i>Lingulella bella</i>
	Tremadocian and older Clarenville Series	Beds with <i>Bryograptus</i> , <i>Shumardia</i> , <i>Parabolina</i> , <i>Princetonia</i> , and <i>Niobe</i> .	Gray and brown shales, with a sandy limestone near the top in at least one locality.	<i>Shumardia</i> <i>Princetonia terranovica</i> <i>Parabolina harrieta</i> <i>Niobe howelli</i> <i>Bellerophon randomi</i> <i>Bryograptus</i>
?	?	<i>Orusia lenticularis</i> Zone	Black and brown shales, and thin sandstones.	<i>Orusia lenticularis</i>
	Upper Cambrian Elliott Cove Series	Zone of <i>Agnostus pisiformis obesus</i> and <i>Olenus</i> .	Black shales and thin gray sandstones.	<i>Agnostus pisiformis obesus</i> <i>Olenus</i>
		<i>Agnostus pisiformis</i> Zone	Dark shales.	<i>Agnostus pisiformis</i>
	?	Age unknown.	Dark sandy shales.	No fossils yet identified.
CAMBRIAN	Middle Cambrian Newfoundland Series	<i>Paradoxides davidis</i> Zone.	Black, brown, and gray shales sometimes with limestone nodules.	<i>Paradoxides davidis</i> <i>Paradoxides rugulosus</i> <i>Corynexochus minor</i> <i>Agnostus punctuosus</i> <i>Agnostus granulatus</i>
		<i>Paradoxides hicksi</i> Zone	Black, brown, and olive shales.	<i>Paradoxides hicksi</i> <i>Agraulos socialis</i> <i>Agnostus fissus</i> <i>Agnostus barrandei</i> <i>Agnostus rex</i> <i>Eodiscus punctatus</i>
		<i>Paradoxides bennetti</i> Zone	Green, and sometimes red, shales with limestone nodules and thin beds of limestone.	<i>Paradoxides bennetti</i> <i>Paradoxides eteminicus</i> <i>Harttella matthewi</i> <i>Conocoryphe elegans</i> <i>Agnostus cf. rex</i>
		<i>Catadoxides</i> Zone.	Green shales, sometimes with manganese shale and limestone in upper part.	<i>Catadoxides magnificus</i>
		<i>Protolenus</i> Zone.	Green and red shales, with some manganese shales and phosphatic beds.	<i>Protolenus</i>
	Lower Cambrian	<i>Callavia</i> Zone, and possibly a pre- <i>Callavia</i> Zone.	Red and green shales, sometimes holding nodules of limestones; red limestones.	<i>Callavia broggeri</i> <i>Strenuella strenua</i> <i>Solenopleura bombifrons</i> <i>Microdiscus bellimarginatus</i> <i>Obolella atlantica</i> <i>Coleoloides typicalis</i>

TABLE II.

CORRELATION OF THE CAMBRIAN-ORDOVICIAN SEDIMENTARY ROCKS OF SOUTHEASTERN NEWFOUNDLAND WITH THOSE OF SOUTHEASTERN CANADA AND NORTHWESTERN EUROPE

SYSTEM	SOUTHEASTERN NEWFOUNDLAND		SOUTHEASTERN CANADA (N.B. and Cape Breton)	NORTHWESTERN EUROPE (Gt. Britain & Scandinavia)	
	Series	Zone			
ORDOVICIAN	Llandeilan?	<i>Lingula fraseri</i> Zone	Not known to be present	Llandeilo of Gt. Britain?	
	Middle and Upper Arenigian, Wabana Series	Beds with <i>Didymograptus</i> and <i>Synhomalonotus</i>	Division 3d of Dr. G. F. Matthew's classification—beds with <i>Didymograptus</i> and <i>Tetragraptus</i>	Middle and Upper Arenig of Great Britain	
	Lower Arenigian, Bell Island Series	Beds with <i>Obolus</i> , <i>Lingulobolus</i> and <i>Sphaerobolus</i>		Lower Arenig of Great Britain	
	Tremadocian and older, Clarendville Series	Beds with <i>Bryograptus</i> , <i>Shumardia</i> , <i>Parabolina</i> , <i>Princetonia</i> and <i>Niobe</i>	Divisions 3a-c of Dr. G. F. Matthew's classification—beds with <i>Ctenopyge</i> , <i>Sphaerophthalmus</i> , <i>Peltura</i> , and <i>Bryograptus</i>	Beds with <i>Shumardia</i> , <i>Parabolina</i> , <i>Peltura</i> , <i>Niobe</i> , and <i>Orusia lenticularis</i>	
?	?	<i>Orusia lenticularis</i> Zone	?		
CAMBRIAN	Upper Cambrian, Elliott Cove Series	Zone of <i>Agnostus pisiformis</i> <i>obesus</i> , and <i>Olenus</i> . <i>Agnostus pisiformis</i> Zone	Division 2 of Dr. G. F. Matthew's classification—beds with varieties of <i>Agnostus pisiformis</i>	Zone of <i>Agnostus pisiformis</i> <i>obesus</i> and <i>Olenus</i> . <i>Agnostus pisiformis</i> Zone (Scandinavia only?)	
	?	Unidentified Zone	<i>Paradoxides forchhammeri</i> Zone?	<i>Paradoxides forchhammeri</i> Zone (Scandinavia only?)	
	Middle Cambrian, Newfoundland Series		<i>Paradoxides davidis</i> Zone	?	<i>Paradoxides davidis</i> Zone
			<i>Paradoxides hicksi</i> Zone	<i>Paradoxides abenacus</i> Zone	<i>Paradoxides hicksi</i> Zone
			<i>Paradoxides bennetti</i> Zone	<i>Paradoxides lamellatus</i> and <i>Paradoxides elemnicus</i> Zones	<i>Paradoxides groomi</i> Zone
	Lower Cambrian		<i>Cataloxides</i> Zone	?	?
			<i>Protolenus</i> Zone	<i>Protolenus</i> Zone	<i>Protolenus</i> Zone
		<i>Callavia</i> Zone, and possibly Pre-Callavia Zone	Etcheminian	<i>Callavia</i> and possibly other Pre-Protolenus Zones	

from the work of the geological survey; 1907.

Jackson, C. T. 1859. (On a *Paradoxides* from St. Mary's Bay, Newfoundland.) Proc. Boston Soc. Nat. Hist., vol. 7, 1859, p. 54.

Jackson, C. T. 1859. (Comparison of specimens of *Paradoxides* from Braintree, Massachusetts, and St. Mary's Bay, Newfoundland.) Proc. Boston Soc. Nat. Hist., vol. 7, 1859, p. 75.

Jukes, J. B. 1839. Report on the geology of Newfoundland, St. John's, Newfoundland, 1839.

Jukes, J. B. 1840. Report on the geology of Newfoundland. Edinburgh. Phil. Jour., New Series, vol. 29, 1840, p. 103-111.

Jukes, J. B. 1840. Report of the progress of the geological survey during 1840. St. John's, Newfoundland, 1840.

Jukes, J. B. 1842. Excursions in and about Newfoundland during the years 1839 and 1840. Two volumes. London, 1842.

Jukes, J. B. 1843. General report of the geological survey of Newfoundland, executed under the direction of the government and legislature of the colony during the years 1839 and 1840. London, 1843.

Logan, W. E. 1865. Geological Survey of Canada. Report of progress from its commencement to 1863, atlas, 1865.

Marcou, J. 1890. The Lower and Middle Taconic of Europe and North America. American Geologist, vol. 5, 1890, p. 356-375, vol. 6, 1890, p. 78-102, 221-233.

Matthew, G. F. 1886. Abstract of a paper on the Cambrian fauna of Cape Breton and Newfoundland. Canadian Record of Science, vol. 2, 1886, No. 4, p. 255-258.

Matthew, G. F. 1887. On the Cambrian faunas of Cape Breton and Newfoundland. Proc. and Trans. of the Roy. Soc. of Canada for 1886, vol. 4, sec. 4, 1887, p. 147-157.

Matthew, G. F. 1896. Faunas of the *Paradoxides* beds in eastern North America, No. 1. Trans. N.Y. Acad. Sci., vol. 15, 1896, p. 192-247.


Matthew, G. F. 1899. A Palaeozoic terrane beneath the Cambrian. Annals N.Y. Acad. Sci., vol. 12, No. 2, 1899, p. 41-56.

Matthew, G. F. 1899. A new Cambrian trilobite. Bull. Nat. Hist. Soc. New Brunswick, No. 17, 1899, p. 136-142.


Matthew, G. F. 1899. Studies on Cambrian faunas, No. 4—Fragments of the Cambrian faunas of Newfoundland. Proc. and Trans. Roy. Soc. Canada for 1899, 2nd ser., vol. 5, sec. 4, No. 3, 1899, p. 67-95.

- Murray, A., and Howley, J. P. 1881. Geological Survey of Newfoundland. London, 1881. (A re-publication of the Reports of Progress of the Geological Survey of Newfoundland for the years 1864 to 1879.)
- Murray, A., and Howley, J. P. 1881. Map of the Peninsula of Avalon, showing distribution of formations. Scale, 4 miles to 1 inch. 1881.
- Rogers, W. B. 1859. (Comparison of *Paradoxides bennetti* with *Paradoxides harlani*.) Proc. Boston Soc. Nat. Hist., vol. 7, 1859, p. 86.
- Salter, J. W. 1859. On the fossils of the *Lingula*-flags or "Zone Primordiale". *Paradoxides* and *Conocephalus* from North America. Quart. Jour. Geol. Soc. London, vol. 15, 1859, p. 551-555.
- van Ingen, G. 1914. Cambrian and Ordovician faunas of southeastern Newfoundland. Bull. Geol. Soc. Amer., vol. 25, 1914, p. 138.
- van Ingen, G. 1914. Table of the Geological Formations of the Cambrian and Ordovician Systems about Conception and Trinity Bays, Newfoundland, and their Northeastern-American and Western-European Equivalents, based upon the 1912-1913 field work. Princeton, 1914.
- Walcott, C. D. 1884. On the Cambrian faunas of North America, preliminary studies. U.S. Geol. Surv., Bull. No. 10, 1884.
- Walcott, C. D. 1888. The stratigraphical succession of the Cambrian Faunas in North America. Nature, vol. 38, 1888, p. 551.
- Walcott, C. D. 1889. Stratigraphic position of the *Olenellus* fauna in North America and Europe. Amer. Jour. of Sci., 3rd ser., vol. 37, 1889, p. 374-392; vol. 38, 1889, p. 29-42.
- Walcott, C. D. 1889. Description of new genera and species of fossils from the Middle Cambrian. Proc. U.S. Nat. Mus., vol. 11, 1889, p. 441-446.
- Walcott, C. D. 1891. The fauna of the Lower Cambrian or *Olenellus* Zone. Tenth Annual Report of the U.S. Geol. Surv., part 1, p. 509-774; 1891. (Report dated 1890, but not issued until 1891.)
- Walcott, C. D. 1891. Correlation Papers—Cambrian. U.S. Geol. Surv., Bull. No. 81, 1891.
- Walcott, C. D. 1891. (The stratigraphic succession of the Cambrian faunas of North America.) Congres geologique international, compte rendu de la 4me session, Londres, 1888, p. 223-225; London, 1891.
- Walcott, C. D. 1891. The North American Continent during Cambrian Time. Twelfth Annual Report of the Director of the U.S. Geol. Surv., part I—Geology, p. 523-568; 1891.
- Walcott, C. D. 1898. Cambrian Brachiopoda; *Obolus* and *Lingulella*, with description of new species. Proc. U.S. Nat. Mus., vol. 21, 1898, p. 385-420.
- Walcott, C. D. 1899. Pre-Cambrian fossiliferous formations. Bull. Geol. Soc. Amer., vol. 10, 1899, p. 199-224.
- Walcott, C. D. 1900. Lower Cambrian terrane in the Atlantic Province. Proc. Wash. Acad. Sci., vol. 1, 1900, p. 301-339.
- Walcott, C. D. 1902. Cambrian Brachiopoda; *Acrotreta*; *Linnarssonella*; *Obolus*; with descriptions of new species. Proc. U.S. Nat. Mus., vol. 25, 1902, p. 577-612.
- Walcott, C. D. 1905. Cambrian Brachiopoda, with descriptions of new genera and species. Proc. U.S. Nat. Mus., vol. 28, 1905, p. 277-337.
- Walcott, C. D. 1910. Cambrian Geology and Palaeontology, No. 6, *Olenellus* and other genera of the *Mesonacidae*. Smithson. Misc. Coll., vol. 53, No. 6, 1910.
- Walcott, C. D. 1912. Cambrian Brachiopoda, U.S. Geol. Surv., Monogr. 51, 1912.
- Walcott, C. D. 1916. Cambrian Geology and Palaeontology, III, No. 3, Cambrian Trilobites. Smithson. Misc. Coll., vol. 64, No. 3, 1916.
- Walcott, C. D. 1916. Cambrian Geology and Palaeontology, III, No. 5, Cambrian Trilobites. Smithson. Misc. Coll., vol. 64, No. 5, 1916.
- Walcott, C. D. 1920. Cambrian Geology and Palaeontology, IV, No. 6—Middle Cambrian Spongiae. Smithson. Misc. Coll., vol. 67, No. 6, p. 261-338; 1920.
- Weston, T. C. 1896. Notes on the geology of Newfoundland. Proceedings and Transactions of the Nova Scotia Institute of Science, vol. 9, 1896, p. 150-157.
- Whiteaves, J. F. 1878. On some Primordial Fossils from Southeastern Newfoundland. Amer. Jour. Sci., 3rd ser., vol. 16, 1878, p. 224-226.
- Willis, B. 1912. Index to the Stratigraphy of North America. U.S. Geol. Surv., Professional Paper 71, 1912.

NOV 16 1926



UNIVERSITY OF ILLINOIS-URBANA



3 0112 098957183



THE GRAPHIC PUBLISHERS
175 Nepean Street
OTTAWA, CANADA